

CONTENTS

Chapter	Title	Page No.
PART I		
PESTICIDE RESIDUE ANALYSIS IN SPICES: METHOD DEVELOPMENT, VALIDATION AND APPLICATION		
CHAPTER 1	INTRODUCTION AND REVIEW	
	Modern pesticides: an overview	1
	Pesticide residues: environmental and health hazards	5
	Analysis of pesticide residues: classical and modern methods	6
	The QuEChERS sample preparation method	8
	Modern adaptations of the QuEChERS method	10
	Instrumentation in pesticide residue analysis	13
	Matrix effects in pesticide residue analysis	16
	Causes of matrix effects in LC-MS/MS	19
	Causes of matrix effects in GC-MS/MS	20
	Correcting matrix effects: LC-MS/MS	22
	Correcting matrix effects: GC-MS/MS	25
	Use of analyte protectants	26
	Residue analytical methods in spices	27
	Analysis of dithiocarbamate residues	28
	Validation of pesticide residue analysis method	29
Scope and objectives of the present investigation	30	
CHAPTER 2	MATERIALS AND METHODS	
	Materials	32
	Instrumentation	32
	Preparation of standard reference solutions	33
	Sample selection and homogenization	32
	General scheme for method development and optimization	34
	QuEChERS sample preparation	37
	Method parameters: GC-M/MS and LC-MS/MS	36
	Preparation of analyte protectant solution for GC-MS/MS	38
	Analysis of pungency and colour in chillies for surrogate matrix studies	38
	Sample preparation for dithiocarbamate (DTC) analysis	39
	Instrumentation for DTC analysis	39
	Method validation	40
	Evaluation of measurement uncertainty	44

CHAPTER 3	RESIDUE ANALYSIS IN SPICES BY UPLC-MS/MS	
	General analytical scheme and establishment of blanks	46
	UPLC-MS/MS method optimization	47
	QuEChERS sample preparation	51
	Optimization of sample: water ratio	51
	Optimization of sample weight	54
	Buffering during extraction step	55
	Optimization of cleanup step	56
	Concentration and reconstitution	60
	Matrix load with optimized cleanup	61
	Evaluation of matrix effects	62
	Method performance	63
	Measurement uncertainty calculation	67
	Conclusions	71
CHAPTER 4	RESIDUE ANALYSIS IN SPICES BY GCMS/MS	
	General analytical scheme and establishment of blanks	73
	GC-MS/MS method optimization	75
	Sample preparation method optimization	76
	Optimization of cleanup conditions	77
	Matrix effects in GC-MS/MS	82
	Method performance	84
	Assessment of measurement uncertainty	86
	Conclusions	89
CHAPTER 5	MITIGATION OF MATRIX EFFECTS IN SPICES	
	Quantitation problems due to matrix effects	90
	Matrix effects in GC-MS/MS	92
	Matrix effects in GC-MS/MS analysis of pesticides	93
	Chemicals as analyte protectants	95
	Matrix effects in LC-MS/MS	101
	Matrix surrogates to mitigate matrix effects in chillies	102
	Study of composition of chilli extracts after cleanup	103
	Use of NVNA as a matrix surrogate for analysis of chilli samples	106
	Application to real samples	110
	Conclusions	114

CHAPTER 6	ANALYSIS OF DITHIOCARBAMATES IN SPICES	
	Structure of DTC compounds	116
	Optimization of instrumental conditions	119
	Method validation	122
	Matrix effects and effects of sample comminution	125
	Safety evaluation of DTC in cardamom and black pepper	127
	Conclusions	128
	REFERENCES	130
PART II		
APPLICATION OF ANALYTICAL METHODS TO REAL SAMPLES		
CHAPTER 1	INTRODUCTION AND REVIEW	
	Risks due to pesticide residues	141
	Fixing MRLs for raw commodities	142
	Dietary exposure and hazard estimation of residues	143
	Fixing MRLs for processed commodities	145
	Dehydration factors in spices	146
	Assessing compliance with MRLs	146
	Scope and objectives of the present study	147
CHAPTER 2	MATERIALS AND METHODS	
	Materials	149
	Instrumentation	149
	Sample collection	150
	Sample preparation and extraction	150
	Instrumental analysis	151
	Assessing compliance with Indian MRLs	151
	Characterization of food safety hazards	152
CHAPTER 3	APPLICATION OF MULTIRESIDUE METHODS TO REAL SAMPLES	
	Residue analysis results	154
	Incidence of residues in spices	156
	Compliance with national MRLs in spices	160
	Food safety risks in spices due to pesticide residues	166
	Hazard characterization of residues in spices	168
	Conclusions	169
	REFERENCES	171
SUMMARY AND RECOMMENDATIONS		174
LIST OF PUBLICATIONS AND CONFERENCE PAPERS		177

LIST OF TABLES

Table No.	Title	Page No.
PART I		
PESTICIDE RESIDUE ANALYSIS IN SPICES: METHOD DEVELOPMENT, VALIDATION AND APPLICATION		
1.1	Classes of synthetic pesticides and mode of functioning	4
1.2	Representative spice matrices with modes of homogenization	34
1.3	Acceptable performance criteria for analytical methods	40
1.4	Optimized UPLC-MS/MS method parameters	47
1.5	Optimized compound-dependent parameters in UPLC-MS/MS	49
1.6	Optimized extraction and QuEChERS cleanup scheme for LC-MS/MS	60
1.7	Linearity equations and correlation coefficient values for pesticides analyzed by LC-MS/MS	64
1.8	Relative standard uncertainty components	69
1.9	Common relative standard uncertainty components – UPLC-MS/MS analysis	70
1.10	Gas chromatographic and mass spectrometric conditions	75
1.11	Optimized MRM transitions in GC-MS/MS	76
1.12	Optimized extraction and QuEChERS cleanup scheme for GC-MS/MS	82
1.13	Validation parameters for target analytes in cumin as a representative matrix	85
1.14	Relative standard uncertainties specific to each analyte compound, at a reference value of 10 $\mu\text{g kg}^{-1}$	86
1.15	Common relative standard uncertainty components: GC-MS/MS analysis	88
1.16	List of analytes and GC-MS/MS retention times	92
1.17	Comparison of matrix effect between extracts of samples MC1 and MC2, with and without dilution	108
1.18	Optimized GC-MS parameter settings for analysis of CS ₂	121
1.19	Accuracy (% recovery), intra-day precision (RSD _r , n = 5) and inter-day precision (RSD _R , n=9) for dithiocarbamates (as CS ₂) in cardamom and black pepper (whole and crushed/ground forms)	123
1.20	Matrix effect (ME, %) in black pepper and cardamom	126
1.21	Safety evaluation of dithiocarbamate residues in cardamom and black pepper	128
PART II		
APPLICATION OF ANALYTICAL METHODS TO REAL SAMPLES		
2.1	Comparison of dietary consumption of cereals and spices	144
2.2	Details of samples collected for survey	150
2.3	Available national MRLs for pesticides for various spices	152
2.4	Classes of pesticides detected in the spice samples studied	167
2.5	Hazard characterization in spices	169

LIST OF FIGURES

Figure No.	Title	Page No.
PART I		
PESTICIDE RESIDUE ANALYSIS IN SPICES: METHOD DEVELOPMENT, VALIDATION AND APPLICATION		
1.1	General chemical structures of commonly used modern pesticides	3
1.2	Steps of the original QuEChERS method	9
1.3	Comparison of the original QuEChERS method with the buffered methods	12
1.4	Instrumentation in tandem mass spectrometry (MS/MS)	14
1.5	Electrospray ionization in LC-MS/MS	15
1.6	Electron ionization in GC-MS/MS	15
1.7	Rotational matrix effects	18
1.8	Schematic representation of matrix effect arising in GC	21
1.9	Strategies for addressing matrix effects	24
1.10	General Scheme for method development and optimization	35
1.11	Optimized UPLC-EI-MS/MS chromatogram	48
1.12	Schematic diagram showing mass spectrometric parameters: LC-MS/MS	49
1.13	Effect of moisture content on extraction efficiency in spices	53
1.14	Effect of sample weight on recovery and precision in cardamom	55
1.15	Optimization of cleanup procedures in spices: UPLC-MS/MS	58
1.16	Matrix load in cleaned extracts: UPLC-MS/MS	61
1.17	Matrix effects (%) of pesticides in four representative spices	66
1.18	Uncertainty components for residue analysis	68
1.19	Expanded uncertainty for 95% confidence limit for UPLC-MS/MS analysis	71
1.20	Optimized chromatogram in GC-MS/MS	77
1.21	Optimization of cleanup procedure in spices: GC-MS/MS	79
1.22	Matrix load in cleaned extracts: GC-MS/MS	81
1.23	Full-scan TIC for extracts of chillies and cardamom, without cleanup (A) and with cleanup (B)	84
1.24	Expanded uncertainty at 95% confidence limit for GC-MS/MS analysis	88
1.25	Schematic representation of quantitation issues due to matrix effects	91
1.26	Comparison of calibration curves in solvent and spice extracts in bifenthrin	94
1.27	GC-MS/MS Matrix effects for pesticides observed in three spices	95
1.28	Chemical structures of analyte protectants used in the study	96
1.29	Effect of volume of AP mix added on peak shapes in 50 mg kg ⁻¹ solvent-based standard in two representative analytes.	97
1.30	Effect of volume of AP mix added on response in 50 mg kg ⁻¹ solvent-based standard as compared to response in matrix matched standards.	99

1.31	Comparison of matrix effects for 50 µg kg ⁻¹ standards in solvent containing 100 µl AP mix /ml of extract and in extracts of three spices	100
1.32	Effect of optimized cleanup step during sample preparation, on A - capsaicinoid content (NHC: nordihydrocapsaicin, CAP: capsaicin, DHC: dihydrocapsaicin) and B - carotenoid content of the chilli extract.	105
1.33	Chemical structures of capsaicinoids in chilli-peppers and synthetic capsaicin: (a) capsaicin, (b) dihydrocapsaicin, (c) nordihydrocapsaicin, (d) homocapsaicin, (e) homodihydrocapsaicin, (f) N-vanillylnonanamide (NVNA, synthetic capsaicin).	106
1.34	Matrix effects of pesticides at concentration of 0.01 mg kg ⁻¹ in (A) surrogate matrix with 50 mg kg ⁻¹ NVNA & MC1 (38,100 SHU) matrix extract diluted to 50%, and (B) surrogate matrix with 50 mg kg ⁻¹ NVNA & MC2 (84,600 SHU) matrix extract diluted to 50%.	109
1.35	Schematic illustration of standard addition technique	111
1.36	Comparison of average values of residue results (µg kg ⁻¹) obtained using 3-point standard addition and surrogate matrix-based calibration (n=3) for three chilli-pepper samples with incurred residues: (A) sample S1 (36,200 SHU), (B) Sample S2 (44,200 SHU) and (C) Sample S3 (58,100 SHU).	113
1.37	Chemical structures of important dithiocarbamate fungicides	117
1.38	Cleavage of thiram to form CS ₂	119
1.39	Optimization of the injection mode in GC-MS. split ratio / split flow	120
1.40	Schematic of the post-run, mid-column backflush option in GC	121
1.41	Chromatogram of CS ₂ in GC-MS, SIM mode	122
1.42	Matrix matched calibration curves for CS ₂ in isooctane, black pepper extract and cardamom extract	125
PART II		
APPLICATION OF ANALYTICAL METHODS TO REAL SAMPLES		
2.1	Using measurement uncertainty in assessing MRL compliance	147
2.2	Incidence of pesticide residues: (A) residues detected in spices (%) out of the 78 compounds tested; (B) overall residue load in the samples of each spice tested	155
2.3	Pesticide residues detected in chillies	156
2.4	Pesticide residues detected in cardamom	157
2.5	Pesticide residues detected in cumin	158
2.6	Pesticide residues detected in ginger	159
2.7	Pesticide residues detected in cinnamon	159
2.8	Pesticide residues detected in curry leaves	160
2.9	Pesticides detected above MRL levels in chillies	161
2.10	Pesticide residues detected above MRL levels in cardamom	162
2.11	Pesticide residues detected above MRL levels in cumin	163
2.12	Pesticide residues detected above MRL levels in ginger	164
2.13	Pesticide residues detected above MRL levels in cinnamon	165
2.14	Pesticide residues detected above MRL levels in curry leaves	166