

REFERENCES

1. Cato MP, Ash HB, Hooper WD. *On Agriculture: Marcus Terentius Varro*. W. Heinemann; 1967.
2. Braidwood RJ. The agricultural revolution. *Scientific American*. 1960;203:130-148.
3. Davis FR. *Banned: A History of Pesticides and the Science of Toxicology*. Yale University Press; 2014.
4. Russell E. *War and Nature: Fighting Humans and Insects with Chemicals from World War I to Silent Spring*. Cambridge University Press; 2001.
5. Oberemok VV, Laikova KV, Gninenko YI, Zaitsev AS, Nyadar PM, Adeyemi TA. A short history of insecticides. *Journal of Plant Protection Research*. 2015;55(3):221-226.
6. Goulson D. Review: An overview of the environmental risks posed by neonicotinoid insecticides. *Journal of Applied Ecology*. 2013;50(4):977-987.
7. Longnecker MP, Rogan WJ, Lucier G. The human health effects of DDT (dichlorodiphenyltrichloroethane) and PCBS (polychlorinated biphenyls) and an overview of organochlorines in public health. *Annual review of public health*. 1997;18(1):211-244.
8. Eskenazi B, Bradman A, Castorina R. Exposures of children to organophosphate pesticides and their potential adverse health effects. *Environmental health perspectives*. 1999;107(suppl 3):409-419.
9. Winter CK. Chronic dietary exposure to pesticide residues in the United States. *International Journal of Food Contamination*. 2015;2(1):1-12.
10. Wesseling C, McConnell R, Partanen T, Hogstedt C. Agricultural pesticide use in developing countries: health effects and research needs. *International journal of health services*. 1997;27(2):273-308.
11. Blacquiere T, Smagghe G, Van Gestel CA, Mommaerts V. Neonicotinoids in bees: a review on concentrations, side-effects and risk assessment. *Ecotoxicology*. 2012;21(4):973-992.
12. Smith TM, Stratton GW. Effects of synthetic pyrethroid insecticides on nontarget organisms. *Residue reviews*. Published online 1986:93-120.
13. Szeto SY, Price PM. Persistence of pesticide residues in mineral and organic soils in the Fraser Valley of British Columbia. *Journal of agricultural and food chemistry*. 1991;39(9):1679-1684.
14. Gevaio B, Semple KT, Jones KC. Bound pesticide residues in soils: a review. *Environmental pollution*. 2000;108(1):3-14.
15. Carson R. *Silent Spring*. Houghton Mifflin Harcourt; 2002.

16. European Union. European Union Maximum Residue Limit Database. Accessed November 17, 2021. <http://ec.europa.eu/food/plant/pesticides/eu-pesticides-database/public/?event=homepage&language=EN>
17. J.F.C.R. Foundation. The Japan Food Chemical Research Foundation Maximum Residue Limit Database. Accessed November 17, 2021. <http://www.m5.ws001.squarestart.ne.jp/foundation/search.html>
18. Codex Alimentarius Commission. Codex MRL Database. Published 2021. Accessed February 1, 2022. <http://www.fao.org/fao-who-codexalimentarius/standards/pesticide-mrls/en/>
19. Food Safety and Standards Authority of India. Food Safety and Standards Regulations. Accessed November 17, 2021. <https://fssai.gov.in/cms/food-safety-and-standards-regulations.php>
20. Mills PA, Onley J, Gaither Ra. Rapid method for chlorinated pesticide residues in nonfatty foods. *Journal of the Association of Official Agricultural Chemists*. 1963;46(2):186-191.
21. Luke MA, Froberg JE, Masumoto HT. Extraction and cleanup of organochlorine, organophosphate, organonitrogen, and hydrocarbon pesticides in produce for determination by gas-liquid chromatography. *Journal of the Association of Official Analytical Chemists*. 1975;58(5):1020-1026.
22. Luke MA, Froberg JE, Doose GM, Masumoto HT. Improved multiresidue gas chromatographic determination of organophosphorus, organonitrogen, and organohalogen pesticides in produce, using flame photometric and electrolytic conductivity detectors. *Journal of the Association of Official Analytical Chemists*. 1981;64(5):1187-1195.
23. Specht W, Pelz S, Gilsbach W. Gas-chromatographic determination of pesticide residues after clean-up by gel-permeation chromatography and mini-silica gel-column chromatography. *Fresenius' journal of analytical chemistry*. 1995;353(2):183-190.
24. Anastassiades M, Scherbaum E. Multiresidue method for determination of pesticide residues in citrus fruits by GC-MSD. *Deutsche Lebensmittel-Rundschau (Germany)*. Published online 1997.
25. Luke M, Cassias I, Yee S. Laboratory Information Bulletin No. 4178. *Office of Regulatory Affairs, US Food and Drug Administration, Rockville, MD*. Published online 1999.
26. Schenck FJ, Callery P, Gannett PM, Daft JR, Lehotay SJ. Comparison of magnesium sulfate and sodium sulfate for removal of water from pesticide extracts of foods. *Journal of AOAC International*. 2002;85(5):1177-1180.
27. Nordmeyer K, Thier HP. Solid-phase extraction for replacing dichloromethane partitioning in pesticide multiresidue analysis. *Zeitschrift für Lebensmitteluntersuchung und-Forschung A*. 1999;208(4):259-263.

28. Casanova JA. Use of solid-phase extraction disks for analysis of moderately polar and nonpolar pesticides in high-moisture foods. *Journal of aoac international*. 1996;79(4):936-940.
29. Richter BE, Jones BA, Ezzell JL, Porter NL, Avdalovic N, Pohl C. Accelerated solvent extraction: a technique for sample preparation. *Analytical chemistry*. 1996;68(6):1033-1039.
30. Barker SA. Matrix solid-phase dispersion. *Journal of chromatography A*. 2000;885(1-2):115-127.
31. Kataoka H, Lord HL, Pawliszyn J. Applications of solid-phase microextraction in food analysis. *Journal of chromatography A*. 2000;880(1-2):35-62.
32. Vas G, Vekey K. Solid-phase microextraction: a powerful sample preparation tool prior to mass spectrometric analysis. *Journal of mass spectrometry*. 2004;39(3):233-254.
33. Pylypiw HM, Arsenault TL, Thetford CM, Incorvia Mattina MJ. Suitability of microwave-assisted extraction for multiresidue pesticide analysis of produce. *Journal of agricultural and food chemistry*. 1997;45(9):3522-3528.
34. Stan HJ. Pesticide residue analysis in foodstuffs applying capillary gas chromatography with mass spectrometric detection: state-of-the-art use of modified DFG-multimethod S19 and automated data evaluation. *Journal of Chromatography A*. 2000;892(1-2):347-377.
35. Anastassiades M, Lehotay SJ, Štajnbaher D, Schenck FJ. Fast and easy multiresidue method employing acetonitrile extraction/partitioning and “dispersive solid-phase extraction” for the determination of pesticide residues in produce. *Journal of AOAC international*. 2003;86(2):412-431.
36. Varela-Martínez DA, González-Sálamo J, González-Curbelo MÁ, Hernández-Borges J. Quick, Easy, Cheap, Effective, Rugged, and Safe (QuEChERS) Extraction. In: *Liquid-Phase Extraction*. Elsevier; 2020:399-437.
37. Lehotay SJ, Kok A de, Hiemstra M, Bodegraven P van. Validation of a fast and easy method for the determination of residues from 229 pesticides in fruits and vegetables using gas and liquid chromatography and mass spectrometric detection. *Journal of AOAC International*. 2005;88(2):595-614.
38. European Committee for Standardization (CEN). Standard Method EN 15662: Food of plant origin - determination of pesticide residues using GC -MS and/or LC-MS/MS following acetonitrile extraction/ partitioning and clean-up by dispersive SPE-QuEChERS method. Published online 2008.
39. Mastovska K, Dorweiler KJ, Lehotay SJ, Wegscheid JS, Szpylka KA. Pesticide multiresidue analysis in cereal grains using modified QuEChERS method combined with automated direct sample introduction GC-TOFMS and UPLC-MS/MS techniques. *Journal of Agricultural and Food Chemistry*. 2010;58(10):5959-5972.

40. González-Curbelo MÁ, Herrera-Herrera AV, Ravelo-Pérez LM, Hernández-Borges J. Sample-preparation methods for pesticide-residue analysis in cereals and derivatives. *TrAC Trends in Analytical Chemistry*. 2012;38:32-51.
41. Li L, Li W, Qin D, Jiang S, Liu F. Application of graphitized carbon black to the QuEChERS method for pesticide multiresidue analysis in spinach. *Journal of AOAC International*. 2009;92(2):538-547.
42. Rajski \Lukasz, Lozano A, Uclés A, Ferrer C, Fernández-Alba AR. Determination of pesticide residues in high oil vegetal commodities by using various multi-residue methods and clean-ups followed by liquid chromatography tandem mass spectrometry. *Journal of Chromatography A*. 2013;1304:109-120.
43. González-Curbelo MÁ, Lehotay SJ, Hernández-Borges J, Rodríguez-Delgado MÁ. Use of ammonium formate in QuEChERS for high-throughput analysis of pesticides in food by fast, low-pressure gas chromatography and liquid chromatography tandem mass spectrometry. *Journal of Chromatography A*. 2014;1358:75-84.
44. Nanita SC, Kaldon LG, Bailey DL. Ammonium salting out extraction with analyte preconcentration for sub-part per billion quantitative analysis in surface, ground and drinking water by flow injection tandem mass spectrometry. *Analytical Methods*. 2015;7(6):2300-2312.
45. Kebarle P, Tang L. From ions in solution to ions in the gas phase-the mechanism of electrospray mass spectrometry. *Analytical chemistry*. 1993;65(22):972A-986A.
46. Kebarle P, Verkerk UH. Electrospray: from ions in solution to ions in the gas phase, what we know now. *Mass spectrometry reviews*. 2009;28(6):898-917.
47. Pal S, Kumar M, Singh R, Kumar N. Evaluation of electron-impact ionization cross sections for molecules. *The Journal of Physical Chemistry A*. 2019;123(19):4314-4321.
48. Märk TD, Dunn GH. *Electron Impact Ionization*. Springer Science & Business Media; 2013.
49. Krueve A, Künnapas A, Herodes K, Leito I. Matrix effects in pesticide multi-residue analysis by liquid chromatography–mass spectrometry. *Journal of chromatography A*. 2008;1187(1-2):58-66.
50. Lehotay SJ, Kwon H, Lucía GA. Sample Preparation and Matrix Effects in the Detection of Chemical Residues in Foods. Accessed December 25, 2020. <http://www.laprw2011.fq.edu.uy/pdf/Lunes/Steven%20Lehotay..pdf>
51. Erney D, Gillespie A, Gilvydis D, Poole C. Explanation of the matrix-induced chromatographic response enhancement of organophosphorus pesticides during open tubular column gas chromatography with splitless or hot on-column injection and flame photometric detection. *Journal of Chromatography A*. 1993;638(1):57-63.
52. Liu C, Dou X, Zhang L, et al. Determination of triazine herbicides and their metabolites in multiple medicinal parts of traditional Chinese medicines using

- streamlined pretreatment and UFLC-ESI-MS/MS. *Chemosphere*. 2018;190:103-113.
53. Chawla S, Patel HK, Gor HN, Vaghela KM, Solanki PP, Shah PG. Evaluation of Matrix Effects in Multiresidue Analysis of Pesticide Residues in Vegetables and Spices by LC-MS/MS. *Journal of AOAC International*. 2017;100(3):616-623. doi:10.5740/jaoacint.17-0048
 54. Furey A, Moriarty M, Bane V, Kinsella B, Lehane M. Ion suppression; a critical review on causes, evaluation, prevention and applications. *Talanta*. 2013;115:104-122.
 55. Antignac JP, de Wasch K, Monteau F, De Brabander H, Andre F, Le Bizec B. The ion suppression phenomenon in liquid chromatography–mass spectrometry and its consequences in the field of residue analysis. *Analytica Chimica Acta*. 2005;529(1-2):129-136.
 56. Matuszewski BK, Constanzer M, Chavez-Eng C. Strategies for the assessment of matrix effect in quantitative bioanalytical methods based on HPLC- MS/MS. *Analytical chemistry*. 2003;75(13):3019-3030.
 57. Mortier KA, Verstraete AG, Zhang GF, Lambert WE. Enhanced method performance due to a shorter chromatographic run-time in a liquid chromatography–tandem mass spectrometry assay for paclitaxel. *Journal of Chromatography A*. 2004;1041(1-2):235-238.
 58. Stüber M, Reemtsma T. Evaluation of three calibration methods to compensate matrix effects in environmental analysis with LC-ESI-MS. *Analytical and Bioanalytical Chemistry*. 2004;378(4):910-916.
 59. Annesley TM. Ion suppression in mass spectrometry. *Clinical chemistry*. 2003;49(7):1041-1044.
 60. Chin C, Zhang Z, Karnes H. A study of matrix effects on an LC/MS/MS assay for olanzapine and desmethyl olanzapine. *Journal of pharmaceutical and biomedical analysis*. 2004;35(5):1149-1167.
 61. Choi B, Hercules D, Gusev A. LC-MS/MS signal suppression effects in the analysis of pesticides in complex environmental matrices. *Fresenius' journal of analytical chemistry*. 2001;369(3):370-377.
 62. Yu S, Xu X min. Study of matrix-induced effects in multi-residue determination of pesticides by online gel permeation chromatography-gas chromatography/mass spectrometry. *Rapid Communications in Mass Spectrometry*. 2012;26(8):963-977.
 63. Rutkowska E, \Lozowicka B, Kaczyński P. Three approaches to minimize matrix effects in residue analysis of multiclass pesticides in dried complex matrices using gas chromatography tandem mass spectrometry. *Food chemistry*. 2019;279:20-29.
 64. Gros M, Petrović M, Barceló D. Development of a multi-residue analytical methodology based on liquid chromatography–tandem mass spectrometry (LC–

MS/MS) for screening and trace level determination of pharmaceuticals in surface and wastewaters. *Talanta*. 2006;70(4):678-690.

65. Andrade J, Terán-Baamonde J, Soto-Ferreiro R, Carlosena A. Interpolation in the standard additions method. *Analytica Chimica Acta*. 2013;780:13-19.
66. Ellison SL, Thompson M. Standard additions: myth and reality. *Analyst*. 2008;133(8):992-997.
67. Ito S, Tsukada K. Matrix effect and correction by standard addition in quantitative liquid chromatographic–mass spectrometric analysis of diarrhetic shellfish poisoning toxins. *Journal of Chromatography A*. 2002;943(1):39-46.
68. Zrostlikova J, Hajšlová J, Poustka J, Begany P. Alternative calibration approaches to compensate the effect of co-extracted matrix components in liquid chromatography–electrospray ionisation tandem mass spectrometry analysis of pesticide residues in plant materials. *Journal of Chromatography A*. 2002;973(1-2):13-26.
69. Larger PJ, Breda M, Fraier D, Hughes H, James CA. Ion-suppression effects in liquid chromatography–tandem mass spectrometry due to a formulation agent, a case study in drug discovery bioanalysis. *Journal of pharmaceutical and biomedical analysis*. 2005;39(1-2):206-216.
70. Villagrasa M, Guillamón M, Eljarrat E, Barceló D. Matrix effect in liquid chromatography–electrospray ionization mass spectrometry analysis of benzoxazinoid derivatives in plant material. *Journal of Chromatography A*. 2007;1157(1-2):108-114.
71. Yang P, Chang JS, Wong JW, et al. Effect of sample dilution on matrix effects in pesticide analysis of several matrices by liquid chromatography–high-resolution mass spectrometry. *Journal of agricultural and food chemistry*. 2015;63(21):5169-5177.
72. Rutkowska E, Łozowicka B, Kaczyński P. Modification of Multiresidue QuEChERS Protocol to Minimize Matrix Effect and Improve Recoveries for Determination of Pesticide Residues in Dried Herbs Followed by GC-MS/MS. *Food Anal Methods*. 2018;11(3):709-724. doi:10.1007/s12161-017-1047-3
73. Kwon H, Lehotay SJ, Geis-Asteggiant L. Variability of matrix effects in liquid and gas chromatography–mass spectrometry analysis of pesticide residues after QuEChERS sample preparation of different food crops. *Journal of chromatography A*. 2012;1270:235-245.
74. Rodríguez-Ramos R, Lehotay SJ, Michlig N, Socas-Rodríguez B, Rodríguez-Delgado MÁ. Critical review and re-assessment of analyte protectants in gas chromatography. *Journal of Chromatography A*. 2020;1632:461596.
75. Maštovská K, Lehotay SJ, Anastassiades M. Combination of analyte protectants to overcome matrix effects in routine GC analysis of pesticide residues in food matrixes. *Analytical Chemistry*. 2005;77(24):8129-8137.

76. Zrostlíková J, Hajšlová J, Godula M, Maštovská K. Performance of programmed temperature vaporizer, pulsed splitless and on-column injection techniques in analysis of pesticide residues in plant matrices. *Journal of Chromatography A*. 2001;937(1-2):73-86.
77. Čajka T, Maštovská K, Lehotay SJ, Hajšlová J. Use of automated direct sample introduction with analyte protectants in the GC–MS analysis of pesticide residues. *Journal of separation science*. 2005;28(9-10):1048-1060.
78. Yarita T, Aoyagi Y, Otake T. Evaluation of the impact of matrix effect on quantification of pesticides in foods by gas chromatography–mass spectrometry using isotope-labeled internal standards. *Journal of Chromatography A*. 2015;1396:109-116.
79. Schenck FJ, Lehotay SJ. Does further clean-up reduce the matrix enhancement effect in gas chromatographic analysis of pesticide residues in food? *Journal of chromatography A*. 2000;868(1):51-61.
80. Yudthavorasit S, Meecharoen W, Leepipatpiboon N. New practical approach for using an analyte protectant for priming in routine gas chromatographic analysis. *Food control*. 2015;48:25-32.
81. Rahman MM, Park JH, Abd El-Aty A, et al. Analysis of kresoxim-methyl and its thermolabile metabolites in Korean plum: An application of pepper leaf matrix as a protectant for GC amenable metabolites. *Journal of separation science*. 2013;36(1):203-211.
82. Poole CF. Matrix-induced response enhancement in pesticide residue analysis by gas chromatography. *Journal of Chromatography A*. 2007;1158(1-2):241-250.
83. Rahman MM, El-Aty AA, Shim JH. Matrix enhancement effect: a blessing or a curse for gas chromatography?—a review. *Analytica chimica acta*. 2013;801:14-21.
84. Erney DR, Poole CF. A study of single compound additives to minimize the matrix induced chromatographic response enhancement observed in the gas chromatography of pesticide residues. *Journal of High Resolution Chromatography*. 1993;16(8):501-503.
85. Erney DR, Pawlowski TM, Poole CF. Matrix-induced peak enhancement of pesticides in gas chromatography: Is there a solution? *Journal of High Resolution Chromatography*. 1997;20(7):375-378.
86. Anastassiades M, Maštovská K, Lehotay SJ. Evaluation of analyte protectants to improve gas chromatographic analysis of pesticides. *Journal of Chromatography A*. 2003;1015(1-2):163-184.
87. FAO/WHO. Codex Committee on Spices and Culinary Herbs (CCSCH) - related standards. Accessed March 27, 2022. <https://www.fao.org/fao-who-codexalimentarius/committees/committee/related-standards/en/?committee=CCSCH>

88. Vázquez PP, Ferrer C, Bueno MM, Fernández-Alba A. Pesticide residues in spices and herbs: Sample preparation methods and determination by chromatographic techniques. *TrAC Trends in Analytical Chemistry*. 2019;115:13-22.
89. Hayward DG, Wong JW. Organohalogen and Organophosphorous Pesticide Method for Ginseng Root □ A Comparison of Gas Chromatography-Single Quadrupole Mass Spectrometry with High Resolution Time-of-Flight Mass Spectrometry. *Analytical chemistry*. 2009;81(14):5716-5723.
90. Chai LK, Elie F. A rapid multi-residue method for pesticide residues determination in white and black pepper (*Piper nigrum* L.). *Food Control*. 2013;32(1):322-326.
91. \Lozowicka B, Jankowska M, Rutkowska E, Hrynko I, Kaczyński P, Miciński J. The evaluation of a fast and simple pesticide multiresidue method in various herbs by gas chromatography. *Journal of Natural Medicines*. 2014;68(1):95-111.
92. Ho YM, Tsoi YK, Leung KSY. Highly sensitive and selective organophosphate screening in twelve commodities of fruits, vegetables and herbal medicines by dispersive liquid–liquid microextraction. *Analytica Chimica Acta*. 2013;775:58-66. doi:<https://doi.org/10.1016/j.aca.2013.02.043>
93. Jadhav M, Thekkumpurath AS, Nakade M, et al. Multiresidue method for targeted screening of pesticide residues in spice cardamom (*Elettaria cardamomum*) by liquid chromatography with tandem mass spectrometry. *Journal of AOAC International*. 2017;100(3):603-609.
94. Shabeer TA, Girame R, Utture S, et al. Optimization of multi-residue method for targeted screening and quantitation of 243 pesticide residues in cardamom (*Elettaria cardamomum*) by gas chromatography tandem mass spectrometry (GC-MS/MS) analysis. *Chemosphere*. 2018;193:447-453.
95. Parmar KD, Kalasariya RL, Litoriya NS, Chawla S, Vaghela KM, Shah PG. Development, validation and evaluation of matrix effect of a QuEChERS-based multiresidue method with low temperature dispersive clean-up for analysis of 104 pesticides in cumin (*Cuminum cyminum*) by LC-MS/MS. *Journal of the Science of Food and Agriculture*. Published online 2021.
96. Girame R, Shabeer TA, Ghosh B, Hingmire S, Natarajan R, Dubey PN. Multi-residue method validation and safety evaluation of pesticide residues in seed spices cumin (*Cuminum cyminum*) and coriander (*Coriandrum sativum*) by gas chromatography tandem mass spectrometry (GC-MS/MS). *Food Chemistry*. 2022;374:131782.
97. Taha SM, Gadalla SA. Development of an efficient method for multi residue analysis of 160 pesticides in herbal plant by ethyl acetate hexane mixture with direct injection to GC-MS/MS. *Talanta*. 2017;174:767-779.
98. Zhang Z, Dong M, Hao X, Han L, Song S, Yao W. Evaluation of cleanup procedures in pesticide multi-residue analysis with QuEChERS in cinnamon bark. *Food chemistry*. 2019;276:140-146.

99. Crnogorac G, Schwack W. Residue analysis of dithiocarbamate fungicides. *TrAC Trends in Analytical Chemistry*. 2009;28(1):40-50.
100. Mujawar S, Utture SC, Fonseca E, Matarrita J, Banerjee K. Validation of a GC–MS method for the estimation of dithiocarbamate fungicide residues and safety evaluation of mancozeb in fruits and vegetables. *Food chemistry*. 2014;150:175-181.
101. Kesari R, Gupta V. A sensitive spectrophotometric method for the determination of dithiocarbamate fungicide and its application in environmental samples. *Talanta*. 1998;45(6):1097-1102.
102. Caldas ED, Conceição MH, Miranda MCC, de Souza LCK, Lima JF. Determination of dithiocarbamate fungicide residues in food by a spectrophotometric method using a vertical disulfide reaction system. *Journal of Agricultural and Food Chemistry*. 2001;49(10):4521-4525.
103. Dubey JK, Heberer T, Stan HJ. Determination of ethylenethiourea in food commodities by a two-step derivatization method and gas chromatography with electron-capture and nitrogen-phosphorus detection. *Journal of Chromatography A*. 1997;765(1):31-38.
104. Gras R, Luong J, Shellie R. Gas chromatography and diode array detection for the direct measurement of carbon disulfide in challenging matrices. *Analytical Methods*. 2017;9(26):3908-3913.
105. Arslan S, Mert ID, Yiğitkaya S, Dagaşan O, Sakallı FN, Oztürk S. The false positive effect of residue of sulphur sources on dithiocarbamate analysis based on CS₂ measurement. *Food Additives & Contaminants: Part A*. 2019;36(1):131-140.
106. Chawla S, Patel H, Kalasariya R, Shah P. Validation and analysis of thiram, a dithiocarbamate, as CS₂ from soybean (*Glycine max*) samples on GC–MS. *International Journal of Environmental Science and Technology*. 2019;16(11):6991-6998.
107. Gustafsson KH, Thompson RA. High-pressure liquid chromatographic determination of fungicidal dithiocarbamates. *Journal of agricultural and food chemistry*. 1981;29(4):729-732.
108. Oellig C, Schwack W. Comparison of HILIC columns for residue analysis of dithiocarbamate fungicides. *Journal of Liquid Chromatography & Related Technologies*. 2017;40(8):415-418.
109. Kakitani A, Yoshioka T, Nagatomi Y, Harayama K. A rapid and sensitive analysis of dithiocarbamate fungicides using modified QuEChERS method and liquid chromatography–tandem mass spectrometry. *Journal of pesticide science*. Published online 2017:D17-025.
110. Xu G, Nie J, Li H, Yan Z, Cheng Y. Determination of mancozeb residue in fruit by derivatization and a modified QuEChERS method using ultraperformance liquid chromatography–tandem mass spectrometry. *Analytical and Bioanalytical Chemistry*. 2017;409(21):5057-5063.

111. Horwitz W. Evaluation of analytical methods used for regulation of foods and drugs. *Analytical chemistry*. 1982;54(1):67-76.
112. European Commission. Guidance document on analytical quality control and method validation procedures for pesticide residue analysis in food and feed (SANTE/12682/2019). Published online January 1, 2020. Accessed March 29, 2020. https://www.eurl-pesticides.eu/docs/public/tmpl_article.asp?CntID=727
113. American Spice Trade Association,. Method 21.3 - Pungency of Capsicums and Their Oleoresins (HPLC Method – Preferred). In: *Official Analytical Methods of the American Spice Trade Association*. ; 2004:111.
114. American Spice Trade Association. Method 20.1 - Extractable Color in Capsicums and Their Oleoresins. In: *The Official Analytical Methods of the American Spice Trade Association*. ; 2004:89.
115. Fu Y, Lu Q, Dou X, Luo J, Yang M, others. Matrix-matched monitoring ion selection strategy for improving the matrix effect and qualitative accuracy in pesticide detection based on UFLC-ESI-MS/MS: A case of Chrysanthemum. *Microchemical Journal*. 2021;160:105681.
116. Ellison SL. Implementing measurement uncertainty for analytical chemistry: the Eurachem Guide for measurement uncertainty. *Metrologia*. 2014;51(4):S199.
117. SANTE/12682. Method Validation & Quality Control Procedures for Pesticide Residues Analysis in Food & Feed (updated 01.01.2020). RR. Published 2019. Accessed December 25, 2020. https://www.eurl-pesticides.eu/docs/public/tmpl_article.asp?CntID=727
118. Grob Jr K, Li Z. Introduction of water and water-containing solvent mixtures in capillary gas chromatography: I. Failure to produce water-wettable precolumns (retention gaps). *Journal of Chromatography A*. 1989;473:381-390.
119. Grob Jr K, Li Z. Introduction of water and water-containing solvent mixtures in capillary gas chromatography: II. Wettability of precolumns by mixtures of organic solvents and water; retention gap techniques. *Journal of Chromatography A*. 1989;473:391-400.
120. Grob Jr K, Neukom H, Li Z. Introduction of water and water-containing solvent mixtures in capillary gas chromatography: III. Water-resistant deactivation of uncoated precolumns? *Journal of Chromatography A*. 1989;473:401-409.
121. Sanatombi K, Sharma G. Capsaicin content and pungency of different Capsicum spp. cultivars. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*. 2008;36(2):89-90.
122. Arimboor R, Natarajan RB, Menon KR, Chandrasekhar LP, Moorkoth V. Red pepper (*Capsicum annum*) carotenoids as a source of natural food colors: analysis and stability—a review. *Journal of food science and technology*. 2015;52(3):1258-1271.
123. Scoville WL. Note on capsicums. *Journal of the American Pharmaceutical Association*. 1912;1(5):453-454.

124. American Spice Trade Association. Method 12.1 - Piperine Content of Black and White Pepper, Their Oleoresins and Soluble Pepper Seasonings. In: *Official Analytical Methods of the American Spice Trade Association.* ; 2018:51.
125. Codex MRL Cardamom. Codex Maximum Residue Limits in HS 0775 - Cardamom. Published 2015. Accessed February 11, 2021. http://www.fao.org/fao-who-codexalimentarius/codex-texts/dbs/pestres/commodities-detail/en/?lang=en&c_id=745
126. Codex MRL Pepper. Codex Maximum Residue Limits in HS 0790 - Pepper, black, white. Published 2015. Accessed February 11, 2021. http://www.fao.org/fao-who-codexalimentarius/codex-texts/dbs/pestres/commodities-detail/en/?c_id=171
127. European Commission. The European Pesticide Database. Published 2021. Accessed February 1, 2022. https://ec.europa.eu/food/plant/pesticides/eu-pesticides-db_en
128. George W, Latimer J. Official method of analysis of AOAC” by AOAC international 21st Edition. Published online 2019.
129. Shabeer TA, Girame R, Utture S, et al. Optimization of multi-residue method for targeted screening and quantitation of 243 pesticide residues in cardamom (*Elettaria cardamomum*) by gas chromatography tandem mass spectrometry (GC-MS/MS) analysis. *Chemosphere.* 2018;193:447-453.
130. Siruguri V, Bhat RV. Assessing intake of spices by pattern of spice use, frequency of consumption and portion size of spices consumed from routinely prepared dishes in southern India. *Nutrition journal.* 2015;14(1):7.
131. Vettorazzi G, Almeida WF, Burin GJ, et al. International safety assessment of pesticides: dithiocarbamate pesticides, ETU, and PTU—a review and update. *Teratogenesis, carcinogenesis, and mutagenesis.* 1995;15(6):313-337.